

# On Unique Predictions for Single Spin Azimuthal Asymmetry

J. P. Ma and Q. Wang

*Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China*

## Abstract

Theoretically there are two approaches to predict single spin azimuthal asymmetries. One is to take transverse momenta of partons into account by using transverse momentum dependent parton distributions, while another is to take asymmetries as a twist-3 effect. The nonperturbative effects in these approaches are parameterized with different matrix elements and predictions can be different. Recently, gauge invariant definitions of transverse momentum dependent parton distributions were derived. With these definitions it can be shown that there are relations between nonperturbative matrix elements in two approaches. These relations may enable us to unify two approaches and to have unique predictions for single spin azimuthal asymmetries. In this letter we derive these relations by using time-reversal symmetry and show that even with these relations the single spin azimuthal asymmetry in Drell-Yan process is predicted differently in different approaches.

PACS numbers: 13.85.Qk, 13.88.+e

Single spin azimuthal asymmetry provides a new tool to study structure of hadrons because the asymmetry is sensitive to correlations between quarks and gluons as partons inside a hadron and to orbital angular momenta of these partons. Experimentally, such an asymmetry was observed for inclusive production of pion in polarized proton antiproton scattering with center-of-mass energy  $\sqrt{s} = 20\text{GeV}$  by E704 collaboration [1]. The asymmetry is large for charged pion, while for  $\pi^0$  production it is consistent with zero when the transverse momentum  $k_T$  is smaller than 3GeV, and it tends to a positive value when  $k_T$  becomes large. In semi inclusive deep-inelastic scattering(SIDIS) significant asymmetries were also observed in production of pion and kaon by HERMES[2]. Asymmetries in polarized proton scattering are currently studied by STAR at RHIC. Large spin effects are observed in their preliminary results after first run. In SIDIS measurements of asymmetries with a transversely polarized target were reported by the SMC collaboration [3]. Experiments with a transversely polarized target are now under study by HERMES and COMPASS [4, 5].

Single spin azimuthal asymmetry is a T-odd effect and helicity-flip amplitudes are involved. Perturbatively T-odd effects can be generated at loop-level in hard scattering of active partons of hadrons. Because the quark-gluon coupling of QCD conserves helicities in the massless limit, the T-odd effects are proportional to quark masses which can be neglected. Therefore the observed T-odd effects can not be explained by those T-odd effects arising from hard scattering and are related to nonperturbative nature of hadrons. Indeed, these T-odd effects can be generated from final- or initial interactions between active partons involved in the hard scattering and remnant partons in hadrons[15, 23, 24]. The effect of these interactions can be represented by gauge links in definitions of parton distributions[23, 24]. Theoretically there are two approaches to explain single spin azimuthal asymmetry by taking nonperturbative nature of hadrons into account. One is to take transverse momenta  $k_T$  of partons in a hadron into account where one uses transverse momentum dependent parton distributions to parameterize nonperturbative effects. For a polarized hadron as an initial state the effect is parameterized by Sivers function[6], while for a hadron observed in a final state the T-odd effect related to this hadron is parameterized by Collins function[7]. For semi-inclusive deep inelastic scattering, both functions can make contributions to the observed single spin azimuthal asymmetry. Single spin azimuthal asymmetry has been studied in terms of these functions [8, 9, 10, 11, 12, 13]. These functions have been also studied with models[14, 15, 16, 17]. Another approach, called Qiu-Sterman mechanism, is that the T-odd effect is produced by taking twist-3 effect into account and it is proportional to quark-gluon correlations inside a hadron[18]. The fact that T-odd effects can be generated at twist-3 level was also pointed out in [19]. This approach was used to make predictions for various processes in [18, 20]. It is interesting to note that at first look the physical reason for the effect is different in different approaches. In the first approach the helicity of a initial hadron is changed because of orbital angular momenta of partons. This can be seen clearly in terms of light-cone wave functions[21]. In the second approach the helicity flip is caused by nonzero spin of the gluon which is correlated with other partons. Predictions based on different approaches are different. A question arises why there are two physical origins for one effect?

This question has been answered partly by recent studies of transverse momentum dependent parton distributions[22, 23, 24], which are involved in the first approach. It has been shown that gauge links in these distributions play an important role to incorporate T-odd effects introduced by final state interactions. In particular, additional gauge links should be included in the definitions of these distributions[24]. With these gauge links it is possible to relate the second  $k_T$  moment of Sivers function to the twist-3 matrix element in the second approach[25]. With relations between nonperturbative matrix elements in different approaches it may be possible to unify two approaches and to have unique predictions for single spin azimuthal asymmetries. In this letter we will show that predictions based on two approaches are still different, although such relations exist. We will show this in detail with Drell-Yan process. Before

showing this we give another derivation of relations between second  $k_T$  moments of T-odd distributions and twist-3 matrix elements by using time-reversal symmetry of QCD.

We consider a proton moving in the  $z$ -direction with the momentum  $P$  and the transverse spin  $\mathbf{s}_T$ . We use a light-cone coordinate system and introduce two light-cone vectors:  $n^\mu = (0, 1, 0, 0)$ ,  $l^\mu = (1, 0, 0, 0)$  and  $n \cdot l = 1$ . Neglecting the proton mass we have  $P^\mu = (P^+, 0, 0, 0)$ . Taking transverse momenta of partons in quark-quark correlation in a proton into account, there are two T-odd parton distribution functions appearing in a Drell-Yan process which take effects of initial state interaction into account. They can be defined as[26]

$$\begin{aligned} f_{1T,DY}^\perp(x, k_T^2) \varepsilon_{\perp\mu\nu} k_T^\mu s_T^\nu &= \frac{1}{4} \int \frac{d\xi^- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \{ \langle P, \mathbf{s}_T | \bar{\psi}(\xi) \gamma^+ V(\xi) \psi(0) | P, \mathbf{s}_T \rangle \\ &\quad - (\mathbf{s}_T \rightarrow -\mathbf{s}_T) \}, \\ h_{1,DY}^\perp(x, k_T^2) k_T^i &= - \int \frac{d\xi^- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \cdot \langle P | \bar{\psi}(\xi) \sigma^{+i} V(\xi) \psi(0) | P \rangle, \end{aligned} \quad (1)$$

with  $\xi^\mu = (0, \xi^-, \xi_T)$  and  $\varepsilon_{\perp\mu\nu} = \varepsilon_{\rho\sigma\mu\nu} n^\rho l^\sigma$ . The momentum  $k$  is  $k^\mu = (xP^+, 0, \mathbf{k}_T)$ . The matrix element in the last line is spin averaged. The function  $f_{1T,DY}^\perp(x, k_T^2)$  is the Sivers function for Drell-Yan processes.  $V(\xi)$  is a product of gauge links to make the matrix element gauge invariant, it takes the effect of initial state interaction in Drell-Yan process into account. If one can take  $V(\xi)$  as a unit matrix, then one can show with time-reversal symmetry that both correlation functions are zero. It is important to note that  $V(\xi)$  is not a unit matrix even in the light-cone gauge  $n \cdot G = 0$ , additional gauge links must be introduced to make the definitions gauge invariant[24]. We will take the light-cone gauge. In this gauge  $V(\xi)$  reads:

$$V(\xi) = V_{-\infty}(\xi_T) = P \exp \left( ig \int_0^{\xi_T} d\xi_T \cdot \mathbf{G}_T(0, \xi^- = -\infty, \xi_T) \right). \quad (2)$$

This gauge link takes effects of initial state interaction into account and it can be derived in a similar way as in SIDIS[24]. The difference is that the gauge link is at  $\xi^- = -\infty$  because it is for initial state interaction.

Under parity- and time-reversal transformation, we obtain for the matrix element:

$$\langle P, \mathbf{s}_T | \bar{\psi}(\xi) \gamma^+ V_{-\infty}(\xi_T) \psi(0) | P, \mathbf{s}_T \rangle = \langle P, -\mathbf{s}_T | \bar{\psi}(\xi) \gamma^+ V_\infty(\xi_T) \psi(0) | P, -\mathbf{s}_T \rangle, \quad (3)$$

with

$$V_\infty(\xi_T) = P \exp \left( ig \int_0^{\xi_T} d\xi_T \cdot \mathbf{G}_T(0, \xi^- = \infty, \xi_T) \right). \quad (4)$$

Similarly one can define two T-odd parton distribution functions appearing in deep inelastic processes which take effects of final state interaction into account. The two functions  $f_{1T,DIS}^\perp(x, k_T^2)$  and  $h_{1,DIS}^\perp(x, k_T^2)$  are defined similarly as in Eq.(1), but with the gauge link  $V(\xi)$  is replaced with  $V_\infty(\xi_T)$ . These two functions are related to those in Drell-Yan processes with time-reversal symmetry. With Eq.( 3) we can write:

$$\begin{aligned} f_{1T,DY}^\perp(x, k_T^2) \varepsilon_{\perp\mu\nu} k_T^\mu s_T^\nu &= \frac{1}{8} \int \frac{d\xi^- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \{ \langle P, \mathbf{s}_T | \bar{\psi}(\xi) \gamma^+ [V_{-\infty}(\xi_T) - V_\infty(\xi_T)] \psi(0) | P, \mathbf{s}_T \rangle \\ &\quad - (\mathbf{s}_T \rightarrow -\mathbf{s}_T) \}, \\ h_{1,DY}^\perp(x, k_T^2) k_T^i &= - \frac{1}{2} \int \frac{d\xi^- d^2\xi_T}{(2\pi)^3} e^{-ik \cdot \xi} \langle P | \bar{\psi}(\xi) \sigma^{+i} [V_{-\infty}(\xi_T) - V_\infty(\xi_T)] \psi(0) | P \rangle. \end{aligned} \quad (5)$$

It is expected that the function  $f_{1T,DY}^\perp$  and  $h_{1,DY}^\perp$  decrease rapidly with increasing  $k_T$ . Then T-odd effects related to them can be estimated at leading order by the second moment of  $k_T$  of the left hand side in Eq.(5):

$$\begin{aligned} K_f^\alpha(x) &= \int d^2 k_T k_T^\alpha f_{1T,DY}^\perp(x, k_T^2) \varepsilon_{\perp\mu\nu} k_T^\mu s_T^\nu = -\frac{1}{2} \varepsilon^{\alpha\sigma} s_{T\sigma} \int d^2 k_T |\mathbf{k}_T|^2 f_{1T,DY}^\perp(x, k_T^2), \\ K_h^{\mu\nu}(x) &= \int d^2 k_T k_T^\mu h_{1,DY}^\perp(x, k_T^2) k_T^\nu = \frac{1}{2} (n^\mu l^\nu + n^\nu l^\mu - g^{\mu\nu}) \int d^2 k_T |\mathbf{k}_T|^2 h_{1,DY}^\perp(x, k_T^2). \end{aligned} \quad (6)$$

With Eq.(5) these moments can be expressed in term of matrix elements. Taking  $K_f^\alpha$  as an example, we have

$$\begin{aligned} K_f^\alpha(x) &= -\frac{1}{8} \int \frac{d\xi^-}{(2\pi)} e^{-ixP^+\xi^-} i \frac{\partial}{\partial \xi_{T\alpha}} \{ \langle P, \mathbf{s}_T | \bar{\psi}(\xi) \gamma^+ [V_{-\infty}(\xi_T) - V_{\infty}(\xi_T)] \psi(0) | P, \mathbf{s}_T \rangle \\ &\quad - (\mathbf{s}_T \rightarrow -\mathbf{s}_T) \} |_{\xi_T=0}. \end{aligned} \quad (7)$$

Carrying out the derivatives we have:

$$\begin{aligned} K_f^\alpha(x) &= \frac{1}{8} \int \frac{d\xi^-}{(2\pi)} e^{-ixP^+\xi^-} \cdot \{ g \langle P, \mathbf{s}_T | \bar{\psi}(\xi^- n) \gamma^+ [G_T^\alpha(0, \infty, 0, 0) - G_T^\alpha(0, -\infty, 0, 0)] \psi(0) | P, \mathbf{s}_T \rangle \\ &\quad - (\mathbf{s}_T \rightarrow -\mathbf{s}_T) \}. \end{aligned} \quad (8)$$

Now one can show that  $K^\alpha$  is related to the twist-3 quark gluon correlation  $T_F(x, x)$  introduced in [18]. The correlation function is defined as:

$$\begin{aligned} T_F(x_1, x_2) \epsilon^{\mu\nu\sigma\rho} n_\nu l_\sigma s_{T\rho} &= -\frac{g}{2} \int \frac{dy_1 dy_2}{4\pi} e^{-iy_2(x_2-x_1)P^+ - iy_1 x_1 P^+} \\ &\quad \cdot \{ \langle P, \mathbf{s}_T | \bar{\psi}(y_1 n) \gamma^+ G^{+\mu}(y_2 n) \psi(0) | P, \mathbf{s}_T \rangle - (\mathbf{s}_T \rightarrow -\mathbf{s}_T) \}. \end{aligned} \quad (9)$$

where we include the coupling constant  $g$  into the definition. It is straightforward to obtain:

$$T_F(x, x) = \int d^2 k_T |\mathbf{k}_T|^2 f_{1T,DY}^\perp(x, k_T^2). \quad (10)$$

Similarly we have:

$$T_H(x, x) = \int d^2 k_T |\mathbf{k}_T|^2 h_{1,DY}^\perp(x, k_T^2), \quad (11)$$

where  $T_H$  is defined with a twist-3 operator:

$$T_H(x_1, x_2) = g \int \frac{dy_1 dy_2}{4\pi} e^{-iy_2(x_2-x_1)P^+ - iy_1 x_1 P^+} \langle P | \bar{\psi}(y_1 n) \sigma^{+\mu} G_\mu^+(y_2 n) \psi(0) | P \rangle. \quad (12)$$

The relations in Eq.(10,11) clearly show that the effect of orbital angular momenta of quarks is closely related to that of quark-gluon correlations because of gauge invariance. These relations also show that the nonperturbative effects in the two approaches for single spin azimuthal asymmetries are the same. However, it should be noted that perturbative coefficients in these two approaches are calculated in different ways. In the first approach one uses  $k_T$ -factorization, while collinear expansion is used in the second approach. If the perturbative coefficients in the two approaches are related in a consistent way so that the predicted single spin azimuthal asymmetries are same, then we may have an unique prediction for single spin azimuthal asymmetries and the question asked before is fully answered. It is difficult to

establish a general relation between the perturbative coefficients, since they are differently calculated in different ways and are different in different processes. But we can show that the single spin azimuthal asymmetry in Drell-Yan process is *differently* predicted between the two approaches.

Now we calculate single spin asymmetry in Drell-Yan process:

$$A(P_A, s_T) + B(P_B) \rightarrow l^-(P_1) + l^+(P_2) + X, \quad (13)$$

where the proton A is transversely polarized with the spin vector  $\mathbf{s}_T$  and moves in the  $+z$ -direction. The  $x$ -direction is chosen as the direction of  $\mathbf{s}_T$ .  $S = (P_A + P_B)^2$ . The hadron B is unpolarized and moves in  $-z$ -direction. We will calculate the single spin azimuthal asymmetry at leading orders, where the lepton pair has a small transverse momentum. We assume that the solid angle  $\Omega(\theta, \phi)$  of the produced lepton in the center-of-mass frame of the produced lepton pair and the invariant mass  $Q^2$  of the lepton pair is observed. The single spin asymmetry is defined as:

$$A_N = \left( \frac{d\sigma(\mathbf{S}_T)}{dQ^2 d\Omega} - \frac{d\sigma(-\mathbf{S}_T)}{dQ^2 d\Omega} \right) / \left( \frac{d\sigma(\mathbf{S}_T)}{dQ^2 d\Omega} + \frac{d\sigma(-\mathbf{S}_T)}{dQ^2 d\Omega} \right). \quad (14)$$

The asymmetry is calculated in [27] with Qiu-Sterman mechanism. The result reads:

$$A_N = -\frac{1}{Q} \cdot \frac{\sin 2\theta \sin \phi}{1 + \cos^2 \theta} \cdot \left( \frac{\sum_q e_q^2 \int dx_A dx_B \delta(Q^2 - x_A x_B S) T_{F,q/A}(x_A, x_A) f_{\bar{q}/B}(x_B)}{\sum_q e_q^2 \int dx_A dx_B f_{q/A}(x_A) f_{\bar{q}/B}(x_B)} + \dots \right) \quad (15)$$

where  $T_{F,q/P}(x, x)$  is defined in Eq.(9). The subscriber  $q/A$  denotes the distribution of  $q$  in hadron A. It should be noted that the summation  $\sum_q$  and also in the below is over all quark and antiquark flavors, i.e.,  $q$  can be an antiquark in the summation. We only keep the term with  $T_{F,q/A}$ . The  $\dots$  represents another term proportional to  $T_{H,\bar{q}/B}$  which is irrelevant in this letter.

In order to make comparison of two approaches we study the single spin asymmetries with  $k_T$  dependent distributions. At tree level the partonic process is just  $q\bar{q} \rightarrow l^+ l^-$ . The cross section can be written:

$$\sigma = e^2 \frac{1}{2S} \int \frac{d^3 P_1}{(2\pi)^3 2P_1^0} \frac{d^3 P_2}{(2\pi)^3 2P_2^0} L_{\mu\nu} W^{\mu\nu} \cdot \frac{1}{Q^4} \quad (16)$$

where the leptonic tensor  $L_{\mu\nu}$  and hadronic tensor  $W^{\mu\nu}$  are given by:

$$\begin{aligned} L^{\mu\nu} &= 4(P_1^\mu P_2^\nu + P_1^\nu P_2^\mu - P_1 \cdot P_2 g^{\mu\nu}), \\ W^{\mu\nu} &= \sum_q e_q^2 \int \frac{d^4 k_A}{(2\pi)^4} \frac{d^4 k_B}{(2\pi)^4} (2\pi)^4 \delta^4(Q - k_A - k_B) \int d^4 \xi_1 d^4 \xi_2 e^{ik_A \cdot \xi_1 + ik_B \cdot \xi_2} \cdot (\gamma^\nu)_{jk} \cdot (\gamma^\mu)_{li} \\ &\quad \cdot [\langle P_A, s_T | \bar{q}_j(0) q_i(\xi_1) | P_A, s_T \rangle \langle P_B | q_k(0) \bar{q}_l(\xi_2) | P_B \rangle + \dots], \end{aligned} \quad (17)$$

where the  $\dots$  denotes power-suppressed terms. The quark  $q$  and  $\bar{q}$  carries the momentum

$$\begin{aligned} k_A &= x_A P_A + k_{AT} \\ k_B &= x_B P_B + k_{BT} \end{aligned} \quad (18)$$

respectively. The quark density matrix with  $k_T$  dependence

$$\Phi_{ij}(x, k_T; P, S) = \frac{1}{2} \int \frac{d\xi^- d^2 \xi_T}{(2\pi)^3} e^{ik \cdot \xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle |_{\xi^+ = 0} \quad (19)$$

can be parameterized as[26]

$$\begin{aligned}\Phi(x, k_T; P, S) &= \frac{1}{4} \{ f_1 \not{l} + f_{1T, DY}^\perp \epsilon_{\mu\nu\rho\sigma} \gamma^\mu l^\nu k_T^\rho s_T^\sigma + g_{1s} \gamma_5 \not{l} + h_{1T} i \sigma_{\mu\nu} \gamma_5 l^\mu s_T^\nu \\ &\quad + h_{1s}^\perp i \sigma_{\mu\nu} \gamma_5 l^\mu k_T^\nu + h_{1, DY}^\perp \sigma_{\mu\nu} k_T^\mu l^\nu \} \end{aligned} \quad (20)$$

where the Sivers function is  $f_{1T, DY}^\perp$ . The function  $f_{1T, DY}^\perp$  and  $h_{1, DY}^\perp$  is defined in Eq.(1). We changed the notation of [26] slightly by replacing  $1/M$  with 1. With this parameterization we have

$$\begin{aligned}W^{\mu\nu} &= \frac{1}{3} \sum_q e_q^2 \int dk_A^+ d^2 k_{AT} dk_B^- d^2 k_{BT} (2\pi)^4 \delta^4(Q - k_A - k_B) (\mathbf{k}_{AT} \times \mathbf{s}_T) \cdot \hat{z} \\ &\quad \cdot f_{1T, q/A}^\perp(k_A) f_{1, \bar{q}/B}(k_B) [g^{\mu\nu} - l^\nu n^\mu - l^\mu n^\nu] + \dots, \end{aligned} \quad (21)$$

where we keep only terms with  $f_{1T, DY}^\perp$ .  $\hat{z}$  denotes the direction of the  $z$ -axis. It should be noted that the total momentum  $Q$  of the lepton pair has nonzero transverse components in general. It depends on transverse momenta of incoming partons. It is now straightforward to calculate the asymmetry defined in Eq.(14). Since the asymmetry is defined as a distribution of variables in the center-mass frame of the lepton pair, we need to specify the frame. We assume that the center-mass frame is obtained from laboratory frame by a Lorentz boost only. This is conveniently used in experiment. In the center-mass frame the lepton  $l^+$  and  $l^-$  has the momentum  $k_2$  and  $k_1$  respectively. The momentum  $k_1$  and  $k_2$  read:

$$\begin{aligned}k_1^\mu &= \frac{\sqrt{Q^2}}{2} (1, \sin \theta \sin \phi, \sin \theta \cos \phi, \cos \phi), \\ k_2^\mu &= \frac{\sqrt{Q^2}}{2} (1, -\sin \theta \sin \phi, -\sin \theta \cos \phi, -\cos \phi) \end{aligned} \quad (22)$$

The momentum  $P_i (i = 1, 2)$  in the laboratory frame is related to  $k_i (i = 1, 2)$  by the boost:

$$\begin{aligned}P_i^0 &= \frac{Q^0}{\sqrt{Q^2}} \left( k_i^0 + \frac{\mathbf{Q} \cdot \mathbf{k}_i}{Q^0} \right), \\ \mathbf{P}_i &= \mathbf{k}_i + \frac{k_i^0}{\sqrt{Q^2}} \mathbf{Q} + \left( \frac{Q^0}{\sqrt{Q^2}} - 1 \right) \frac{\mathbf{Q} \cdot \mathbf{k}_i}{\mathbf{Q} \cdot \mathbf{Q}} \mathbf{Q}. \end{aligned} \quad (23)$$

The phase space integration is invariant under the boost. Using Eq.(23) one can express  $L^{\mu\nu}$  in term of  $k_1, k_2$  and  $Q$ . For transverse momentum independent parton distributions one expects in general that they decrease rapidly with increasing transverse momenta. Hence an expansion of the perturbative part in transverse momenta is an good approximation. Keeping the first non-zero order in the expansion of  $k_{AT}$  and  $k_{BT}$ , we obtain the asymmetry:

$$\begin{aligned}A_N &= \frac{1}{Q} \cdot \frac{\sin 2\theta \sin \phi}{1 + \cos^2 \theta} \sum_q e_q^2 \int dx_A dx_B \delta(Q^2 - x_A x_B S) \frac{x_B - x_A}{2(\sqrt{x_A} + \sqrt{x_B})^2} \\ &\quad \cdot f_{\bar{q}/B}(x_B) \cdot \int d^2 k_T |\mathbf{k}_T|^2 f_{1T, DY, q/A}^\perp(x_A, k_T^2) / \sum_q e_q^2 \int dx_A dx_B f_{q/A}(x_A) f_{\bar{q}/B}(x_B) + \dots \\ &= -\frac{1}{Q} \cdot \frac{\sin 2\theta \sin \phi}{1 + \cos^2 \theta} \sum_q e_q^2 \int dx_A dx_B \delta(Q^2 - x_A x_B S) \frac{x_A - x_B}{2(\sqrt{x_A} + \sqrt{x_B})^2} \\ &\quad \cdot f_{\bar{q}/B}(x_B) \cdot T_{F, q/P}(x_A, x_A) / \sum_q e_q^2 \int dx_A dx_B f_{q/A}(x_A) f_{\bar{q}/B}(x_B) + \dots \end{aligned} \quad (24)$$

In the above equation we have assumed that the initial hadrons are in a center-mass frame, i.e.,  $P_A^0 = P_B^0$ . In the last step we have used the relation in Eq.(10). Again, the summation  $\sum_q$  is over all quark and antiquark flavors. It is clearly that the asymmetry here is different than that in Eq.(15) because of the factor  $(x_A - x_B)/2(\sqrt{x_A} + \sqrt{x_B})^2$ . If the factor was 1, then the asymmetry would be the same. Hence, the asymmetry obtained by two approaches will have the same angular distribution but the normalization is different. Since the factor can be positive or negative, the asymmetry from the two approaches can even have different signs.

It should be noted that the hadronic tensor calculated with the parameterization in Eq.(20) is not invariant under electromagnetic gauge transformation. This can be seen by evaluating  $Q_\mu W^{\mu\nu}$  with  $W^{\mu\nu}$  given in Eq.(21). The reason is that the partons involved in the hard scattering have nonzero transverse momenta and  $\gamma \cdot l = \gamma^-$  is contracted with the hard part of  $q\bar{q} \rightarrow l^+l^-$  according to the first two terms in Eq.(20). With nonzero transverse momenta the contraction with  $\gamma^-$  does not make the  $q\bar{q}$  pair on-shell. Hence the  $U_{em}(1)$  gauge invariance is not preserved. It is also indicated[28] that the decomposition in Eq.(20) may need to be reexamined because the density matrix element also depends on the vector  $n^\mu$  due to gauge links. This dependence is neglected in Eq.(20). There are possibly many ways to restore the gauge invariance. In this letter we simply make the initial parton on-shell by replacing  $\gamma^-$  with  $\gamma \cdot k/n \cdot k$  for the first two terms in Eq.(20), i.e.,

$$\Phi(x, k_T; P, s_T) = \frac{1}{4} \{ f_1(x, k_T^2) \frac{\gamma \cdot k}{k \cdot n} + f_{1T, DY}^\perp(x, k_T^2) \frac{\gamma \cdot k}{k \cdot n} \varepsilon_{\perp \rho\sigma} k_T^\rho s_T^\sigma + \dots \}. \quad (25)$$

The hadronic tensor obtained with Eq.(25) is  $U_{em}(1)$ -gauge invariant. It is:

$$W^{\mu\nu} = \frac{1}{3} \sum_q e_q^2 \int dk_A^+ d^2 k_{AT} dk_B^- d^2 k_{BT} (2\pi)^4 \delta^4(Q - k_A - k_B) (\mathbf{k}_{AT} \times \mathbf{s}_T) \cdot \hat{z} \\ \frac{1}{(k_A \cdot n)(k_B \cdot l)} [g^{\mu\nu} k_A \cdot k_B - k_A^\mu k_B^\nu - k_A^\nu k_B^\mu] f_{1T, DY, q/A}^\perp(k_A) f_{1, \bar{q}/B}(k_B). \quad (26)$$

It is straightforward to show  $Q_\mu W^{\mu\nu} \sim k_A^2 k_B^\nu + k_B^2 k_A^\nu \sim k_T^2$ . Hence the gauge invariance is preserved up to order  $k_T^2$ . The asymmetry calculated with this tensor will be gauge invariant. The result of  $A_N$  can be obtained from Eq.(24) by replacing the factor  $(x_B - x_A)/2(\sqrt{x_A} + \sqrt{x_B})^2$  with  $\sqrt{x_B}/(\sqrt{x_A} + \sqrt{x_B})$ . Therefore, even after we make the hadronic tensor gauge invariant, the obtained asymmetry  $A_N$  is still different than that in Eq.(15) from the second approach. It is interesting to see how the same asymmetry in Eq.(15) can be obtained by starting from Eq.(26). If we replace the tensor  $[g^{\mu\nu} k_A \cdot k_B - k_A^\mu k_B^\nu - k_A^\nu k_B^\mu]$  with  $[g^{\mu\nu} k_A \cdot k'_B - k_A^\mu k_B'^\nu - k_A^\nu k_B'^\mu]$  where  $k_B'^\mu = (0, k_B^-, -\mathbf{k}_{AT})$  and neglect the dependence of lepton momenta on transverse momenta of incoming partons, we indeed obtain the asymmetry in Eq.(15) with the same normalization, but with an extra negative sign. However, the transverse momentum of the lepton pair can not be neglected and we can not do the replacement.

To summarize: There are two different approaches for single spin azimuthal asymmetries. Using time-reversal symmetry, we give in this letter an detailed derivation of the relations between  $k_T$  dependent T-odd distributions and twist-3 quark-gluon correlators, which are used in different approaches, respectively. These relations show that the physical origin in the two different approaches for single spin azimuthal asymmetries is the same because of the gauge invariance. With these relations it may be expected to unify these two approaches and to delivery an unique prediction for single spin azimuthal asymmetries. We have studied in detail the single spin azimuthal asymmetry in Drell-Yan process with Sivers function and found that predictions from different approaches are different even with these relations. The  $k_T$  factorization used for single spin azimuthal asymmetries does not respect the  $U_{em}(1)$ -gauge invariance.

This problem may be solved by changing the projection of the perturbative part slightly. But even after this changing the predicted asymmetry is still different. Our study shows clearly that different approaches give different predictions for the same physical effect in Drell-Yan process and one can expect that the same situation will also appear in other processes. Therefore, at present we have not an unique prediction for single spin azimuthal asymmetries in Drell-Yan process at least and this problem needs to be studied further.

### **Acknowledgements**

We would like to thank Prof. X.D. Ji and Prof. J.W. Qiu for useful discussions. This work is supported by National Nature Science Foundation of P. R. China.



## References

- [1] E581 and E704 Collaboration, Phys. Lett. B261 (1991)201;  
E704 Collaboration, Phys. Lett. B264(1991)462;  
E704 Collaboration, Phys. Rev. D53 (1996)4747.
- [2] H. Avakian, HERMES Collaboration, Nucl. Phys. Proc. Suppl. **79** (1999) 523;  
A. Airapetian et al. HERMES Collaboration, Phys. Rev. Lett. **84** (2000) 4047;  
A. Airapetian et al. HERMES Collaboration, Phys. Rev. D64 (2001) 097101;  
A. Airapetian et al. HERMES Collaboration, Phys. Lett. B562 (2003) 182.
- [3] A. Bravar, Spin Muon Collaboration, Nucl. Phys. Proc. Suppl. **79**(1999)520.
- [4] N. C. Makins and M. Duren, HERMES Collaboration, Acta Phys. Polon. B33 (2002)3737;  
N. C. Makins, HERMES Collaboration, Nucl. Phys. A711 (2002) 41;
- [5] J. M. Le Goff, COMPASS Collaboration, Nucl. Phys. A711 (2002) 56;  
M. Lamanna, COMPASS Collaboration, Nucl. Phys. A711 (2002) 50.
- [6] D. Sivers, Phys. Rev. D41 (1990) 83, Phys. Rev. D43 (1991) 261.
- [7] J. C. Collins, Nucl. Phys. B396 (1993) 161.
- [8] M. Anselmino, M. Boglione and F. Murgia, Phys. Lett. B362 (1995) 164;  
M. Anselmino and F. Murgia, Phys. Lett. B442 (1998) 470;  
M. Anselmino and F. Murgia, Phys. Lett. B483 (2000) 74;  
M. Anselmino, U. D'Alesio and F. Murgia, Phys.Rev. D67 (2003) 074010.
- [9] P. J. Mulders and R. D. Tangerman, Nucl. Phys. B461 (1996) 197 [Erratum-ibid. B484 (1997) 538];  
D. Boer, Phys. Rev. D60 (1999)014012.
- [10] E. De Sanctis, W. D. Nowak and K. A. Oganessian, Phys. Lett. B483 (2000) 69;  
V. A. Korotkov, W. D. Nowak and K. A. Oganessian, Eur. Phys. J. C18 (2001) 639;  
K. A. Oganessian, N. Bianchi, E. De Sanctis and W. D. Nowak, Nucl. Phys. A689 (2001) 784;
- [11] A. V. Efremov, K. Goeke, M. V. Polyakov and D. Urbano, Phys. Lett. B478 (2000) 94;  
A. V. Efremov, K. Goeke and P. Schweitzer, Eur. Phys. J. C24 (2002) 407, Nucl. Phys. A711 (2002) 84, Phys. Lett. B522 (2001) 37, [Erratum-ibid. B544 (2002) 389], Phys. Lett. B568 (2003) 63.
- [12] B. Q. Ma, I. Schmidt and J. J. Yang, Phys. Rev. D66 (2002) 094001, Phys. Rev. D65 (2002) 034010, Phys. Rev. D63 (2001) 037501.
- [13] Z. T. Liang and T. C. Meng, Z.Phys. A344 (1992)171;  
C.Boros, Z. T. Liang and T. C. Meng, Phys.Rev.Lett. **70** (1993) 1751.

- [14] P. V. Pobylitsa and M. V. Polyakov, Phys. Lett. B389 (1996) 350,  
P. Schweitzer *et al.*, Phys. Rev. D64 (2001) 034013,  
A. Bacchetta *et al.*, Phys.Lett. B506 (2001) 155.
- [15] S. J. Brodsky, D. S. Hwang and I. Schmidt, Phys. Lett. B530 (2002) 99,  
D. Boer, S. J. Brodsky and D. S. Hwang, Phys. Rev. D67 (2003) 054003,  
S. J. Brodsky, D. S. Hwang and I. Schmidt, Phys. Lett. B553,
- [16] L. P. Gamberg, G. R. Goldstein and K. A. Oganessyan, arXiv:hep-ph/0301018.
- [17] F. Yuan, arXiv:hep-ph/0308157,  
A. Bacchetta, A. Schäfer and J.J. Yang, arXiv: hep-ph/0309246.
- [18] J. W. Qiu and G. Sterman, Phys. Rev. Lett **67** (1991) 2264, Nucl. Phys. B378 (1992) 52, Phys. Rev. D59 (1998) 014004.
- [19] A.V. Efremov and O.V. Teryaev, Sov. J. Nucl. Phys. **36** 1982 1, Phys. Lett. B150 (1985) 383.
- [20] Y. Kanazawa and Y. Koike, Phys. Lett. B478 (2000) 121, *ibid.* B490 (2000) 99, Phys. Rev. D64 (2001) 034019.
- [21] X. D. Ji, J. P. Ma and F. Yuan, Nucl. Phys. B652 (2003) 383.
- [22] S. J. Brodsky et al., Phys. Rev. D65 (2002) 114025.
- [23] J. C. Collins, Phys. Lett. B536 (2002) 43.
- [24] X. D. Ji and F. Yuan, Phys. Lett. B543 (2002) 66,  
A. V. Belitsky, X. D. Ji and F. Yuan, Nucl. Phys. B656 (2003) 165.
- [25] D. Boer, P. J. Mulders and F. Pijlman, Nucl. Phys. B667 (2003) 201.
- [26] D. Boer and P. J. Mulders, Phys. Rev. D57 (1998) 5780.
- [27] N. Hammon, O. Teryaev, and A. Schafer, Phys. Lett. B390 (1997) 409,  
D. Boer, P. J. Mulders and O. V. Teryaev, Phys. Rev. D57 (1998) 3057,  
D. Boer and P. J. Mulders, Nucl. Phys. B569 (2000) 505,  
D. Boer and J. W. Qiu, Phys. Rev. D65 (2002) 034008.
- [28] K. Goeke, A. Metz, P. V. Pobylitsa, M. V. Polyakov, Phys. Lett. B567 (2003) 27.